

NOVARTIS AG
Form 6-K
October 03, 2005

SECURITIES AND EXCHANGE COMMISSION

Washington, D.C. 20549

FORM 6-K

REPORT OF FOREIGN PRIVATE ISSUER
PURSUANT TO RULE 13a-16 or 15d-16 OF
THE SECURITIES EXCHANGE ACT OF 1934

Report on Form 6-K for September 2005

(Commission File No. 1-15024)

Novartis AG

(Name of Registrant)

Lichtstrasse 35

4056 Basel

Switzerland

(Address of Principal Executive Offices)

Indicate by check mark whether the registrant files or will file annual reports under cover of Form 20-F or Form 40-F:

Edgar Filing: NOVARTIS AG - Form 6-K

Form 20-F: Form 40-F:

Indicate by check mark if the registrant is submitting the Form 6-K in paper as permitted by Regulation S-T Rule 101(b)(1):

Yes: No:

Indicate by check mark if the registrant is submitting the Form 6-K in paper as permitted by Regulation S-T Rule 101(b)(7):

Yes: No:

Indicate by check mark whether the registrant by furnishing the information contained in this form is also thereby furnishing the information to the Commission pursuant to Rule 12g3-2(b) under the Securities Exchange Act of 1934.

Yes: No:

Enclosures:

1. FDA advisory committee votes unanimously to recommend approval of Exjade® for once-daily oral treatment of chronic iron overload due to blood transfusions (Basel, September 30, 2005)

2. New data show that Novartis indacaterol provides 24-hour efficacy with single dose in patients with asthma and COPD
(Copenhagen, September 19, 2005)

3. Three-year study confirms long-term benefits of Xolair® for patients with severe allergic asthma
(Basel, September 19, 2005)

4. Novartis signs global licensing agreement to develop new topical applications of terbinafine the most prescribed oral treatment for fungal nail infections (Basel, September 15, 2005)

5. Sandoz Files Lawsuit Seeking FDA Ruling on Omnitrope
(Holzkirchen, Germany, September 13, 2005)

6. Novartis and Alnylam create major alliance to discover RNAi therapeutics
(Basel, Switzerland, and Cambridge, MA, September 7, 2005)

7. New data demonstrate benefits of Diovan® in reducing life-threatening complications from narrowed arteries after a heart attack (Stockholm, September 5, 2005)

Novartis International AG

Novartis Communications

CH-4002 Basel

Switzerland

Tel +41 61 324 2200

Fax + 41 61 324 3300

Internet Address:

<http://www.novartis.com>

MEDIA RELEASE

COMMUNIQUE AUX MEDIAS

MEDIENMITTEILUNG

FDA advisory committee votes unanimously to recommend approval of Exjade® for once-daily oral treatment of chronic iron overload due to blood transfusions

Novel, once-daily oral iron chelator may eliminate need for lengthy, burdensome infusion with current standard of care

Committee supports use of Exjade in patients age 6 and older

Review is based on data from largest global clinical trials program of its type, including 1,000+ adults and children

Basel, September 30, 2005 Novartis announced today that the Blood Products Advisory Committee (BPAC) of the U.S. Food and Drug Administration (FDA) gave a positive review of Exjade® (deferasirox).

Exjade is currently under priority review by the FDA as the first and only once-daily oral iron chelator for the treatment of chronic iron overload due to blood transfusions in adults and children. A newly designed molecule, Exjade is administered as a once-daily drink after tablet(s) are dispersed in a glass of water or orange juice.

The committee unanimously voted to recommend approval for use of Exjade for patients with chronic iron overload due to blood transfusions. The FDA generally follows the recommendations of its advisory committees, although it is not obligated to do so. If approved, Exjade may be a major advance in iron chelation therapy in the U.S.

We believe the Blood Products Advisory Committee has recognized the potential of Exjade to dramatically improve the management of patients with transfusional iron overload, said Diane Young, MD, vice president, global head, Clinical Development, Novartis Oncology. During the next several weeks we will work closely with the FDA to answer any remaining questions, in hopes of bringing this new treatment option to patients

as soon as possible.

Exjade, which has been designated an orphan drug in both the EU and the US, is currently under priority review in the US, Canada, Switzerland, Australia and New Zealand. Additional regulatory submissions have been filed around the world.

Iron overload is a cumulative, potentially life-threatening, unavoidable consequence of frequent blood transfusions used to treat certain rare, chronic blood disorders, including thalassemia, sickle cell disease, other rare anemias and myelodysplastic syndromes. Signs of iron overload may be

detected after 10 to 20 blood transfusions. If left undiagnosed or untreated, the excess iron in the body leads to damage to the liver, heart and endocrine glands. Iron chelation has been demonstrated to be the only effective treatment for transfusion-related iron overload.

Filing Data

The Exjade filings were based on the results of a pivotal clinical trials program, including a Phase III head-to-head trial vs. Desferal® (deferrioxamine), which showed that Exjade significantly reduced liver iron concentration (LIC) at doses of 20-30 mg/kg/day. These clinical trials, which included more than 1,000 adults and children, were part of the largest prospective global clinical trials program ever implemented for an investigational iron chelator. LIC is the accepted indicator for body iron content in patients receiving blood transfusions. The studies demonstrated that Exjade at 20-30 mg/kg/day led to the maintenance or reduction of iron burden in transfused patients with thalassemia, sickle cell disease, other rare anemias and myelodysplastic syndromes. In the clinical studies, Exjade was generally well tolerated, with the most frequently reported adverse events being nausea, vomiting, diarrhea, abdominal pain, skin rash and increases in serum creatinine. As with deferrioxamine, cases of ocular and auditory disturbances have been reported.

About Iron Chelation

In iron chelation an agent binds to iron in the body and helps remove it through the urine and/or feces. To date, only deferrioxamine is globally available for the first-line treatment of transfusional iron overload. While deferrioxamine is effective, it typically requires subcutaneous infusion lasting eight to twelve hours per day, for five to seven days a week for as long as the patient continues to receive blood transfusions or has excess iron within their body. In many patients the need for transfusions may be life-long. However, due to the inconvenience and discomfort associated with the administration of deferrioxamine, many patients choose not to undergo iron chelation therapy, exposing themselves to the dangers of iron overload. Novartis believes the approval of Exjade would, therefore, not only help patients currently receiving iron chelation, but also extend the benefits of iron chelation to those not currently undergoing therapy.

Additional Information

Clinical trials with Exjade are ongoing. To learn more about Exjade clinical trials, health care providers can call either 800 561 1376 in the US, or +44 (0) 1506 814899 outside the US.

The foregoing release contains forward-looking statements that can be identified by terminology such as if approved, Exjade may be , believes, potential, will, in hopes of bringing, would, dramatically improve, recommended, or similar expressions, or by express or implied discussion regarding potential marketing approvals or future sales of Exjade. Such forward-looking statements involve known and unknown risks, uncertainties and other factors that may cause actual results with Exjade to be materially different from any future results, performance or achievements expressed or implied by such statements. There can be no guarantee that Exjade will receive any marketing approvals, or that it will reach any particular sales levels. In particular, management's expectations regarding commercialization of Exjade could be affected by, among other things, additional analysis of Exjade clinical data; new clinical data; unexpected clinical trial results; unexpected regulatory actions or delays or government regulation generally; the company's ability to obtain or maintain patent or other proprietary intellectual property protection; competition in general; increased government, industry, and general public pricing pressures; and other risks and factors referred to in Novartis AG's current Form 20-F on file with the U.S. Securities and Exchange Commission. Should one or more of these risks or uncertainties materialize, or should underlying assumptions prove incorrect, actual results may vary materially from those anticipated, believed, estimated or expected. Novartis is providing the information in this press release as of this date and does not undertake any obligation to update any forward-looking

statements contained in this press release as a result of new information, future events or otherwise.

For prescribing information on Desferal (deferoxamine) please contact your local Novartis affiliate.

About Novartis

Novartis AG (NYSE: NVS) is a world leader in pharmaceuticals and consumer health. In 2004, the Novartis group of companies' businesses achieved sales of USD 28.2 billion and a pro forma net income of USD 5.8 billion. The group invested approximately USD 4.2 billion in R&D. Headquartered in Basel, Switzerland, Novartis group companies employ approximately 83,700 people and operate in over 140 countries around the world. For further information, please consult <http://www.novartis.com>.

###

© 2001 Novartis AG, Legal Disclaimer

Novartis International AG
Novartis Communications
CH-4002 Basel
Switzerland

Tel +41 61 324 2200
Fax + 41 61 324 3300
Internet Address:
<http://www.novartis.com>

MEDIA RELEASE COMMUNIQUE AUX MEDIAS MEDIENMITTEILUNG

New data show that Novartis indacaterol provides 24-hour efficacy with single dose in patients with asthma and COPD

Studies confirm indacaterol could become first in new class of once-daily beta2-agonists

Copenhagen, September 19, 2005 Novartis announced today that its development compound indacaterol demonstrated effective and well-tolerated treatment of asthma and chronic obstructive pulmonary disease (COPD) over 24 hours with a rapid onset of action, according to new data presented at the congress of the European Respiratory Society (ERS) in Copenhagen, Denmark.

The combination of 24-hour efficacy and a reassuring safety profile suggests that in time, a once-daily dose of indacaterol could become a new standard of care for bronchodilation in asthma and COPD, said Joerg Reinhardt, Global Head of Development, Novartis Pharma AG. We are now concentrating on the development of this important new therapy for the benefit of patients who suffer from these debilitating and sometimes fatal diseases.

The results of a series of clinical studies presented at the meeting show that indacaterol (formerly known as QAB149) could become the first in a new class of once-daily beta2-agonists, offering potential therapeutic benefits for patients with asthma and COPD. Indacaterol could be the first beta2-agonist to be taken only once-daily providing full 24 hour symptom control with a single administration, in contrast to currently-available long-acting beta2-agonists (LABAs) such as salmeterol and formoterol which have to be taken twice-daily.

Key indacaterol data presented at ERS included those demonstrating its 24-hour bronchodilator efficacy in COPD, as well as in asthma. Even at high doses, indacaterol demonstrated a good overall safety profile with no concerns over key adverse events sometimes associated with beta2-agonists. These data reinforce the results of preclinical studies.

Considered together, these results provide important insights into the future therapeutic potential of indacaterol, the first in a new generation of drugs that could accurately be described as once-daily beta2-agonists, said Prof. Stephen Holgate, Southampton General Hospital, UK. For patients with asthma or COPD, indacaterol could provide important clinical benefits in terms of improved compliance and more rapid and reliable long-term control of the potentially life-threatening symptoms of breathlessness and bronchial constriction associated with these

conditions.

24-hour efficacy in asthma and COPD

The efficacy of indacaterol in both asthma and COPD was demonstrated in a series of placebo-controlled

clinical studies using once-daily doses of indacaterol ranging from 25 to 2000 µg.(1)-(7) The duration of action of indacaterol was found to be largely independent of dose, with superior bronchodilation to placebo demonstrated at 24 hours after a single dose.(1)

The efficacy of indacaterol in patients with asthma was further investigated in three multiple-dose studies of 7, 14 and 28 days duration.(3)-(5) In these studies, the 24-hour bronchodilator efficacy of indacaterol observed on the first day was maintained for the duration of the studies, suggesting that regular use of indacaterol is not associated with the development of tolerance, or tachyphylaxis. Indacaterol also demonstrated 24-hour bronchodilator efficacy with no evidence of tachyphylaxis in patients with COPD.(6),(7)

Strong safety profile

Data presented at ERS demonstrate that the 24-hour efficacy of indacaterol in asthma is accompanied by a positive safety profile. Single indacaterol doses up to 2000 µg were well-tolerated and were not associated with any clinically significant changes in known class effect adverse events such as hypokalaemia, hyperglycaemia, increased heart rate or altered QTc interval.(1),(2)

These single-dose results were confirmed in multiple-dose asthma studies, in which indacaterol doses up to 800 µg once-daily for up to 28 days were associated with a good cardiovascular safety profile and no clinically relevant effects on blood pressure, QTc, glucose or potassium levels.(3)-(5)

The positive safety profile of indacaterol in asthma was also observed in patients with COPD, at doses up to 800 µg for up to 28 days.(6),(7)

Findings supported by preclinical data

Data from preclinical studies were presented at ERS and confirmed the results of clinical studies, with indacaterol acting as a potent beta2-agonist and displaying high intrinsic efficacy in isolated human bronchial tissue.(8) In other studies, indacaterol demonstrated a sustained duration of action in isolated human bronchial tissue(9) and in guinea pigs.(10) Preclinical data also support the lack of tachyphylaxis observed in clinical studies.(11)

The safety of indacaterol was also examined in the preclinical development programme. In a series of *in vitro* and *in vivo* studies, the safety of indacaterol was compared with that of two long-acting beta2-agonists (formoterol and salmeterol) in doses providing equivalent degrees of bronchodilation.(12) Indacaterol demonstrated a better cardiovascular safety profile than formoterol and salmeterol.(12) Importantly, studies using isolated human bronchial tissue suggest that indacaterol will not antagonise the bronchorelaxant effect of short-acting beta2-agonists, and therefore should not interfere with rescue medication use.(8)

About indacaterol

Edgar Filing: NOVARTIS AG - Form 6-K

Indacaterol (previously known as QAB149) works by stimulating beta2-receptors in the smooth muscle of the airways. This causes relaxation of the muscle, thereby increasing the diameter of the airways, which become constricted in asthma and COPD.

Indacaterol is being developed both as monotherapy and as a fixed-dose combination with drugs such as NVA237, a long-acting anti-muscarinic agent for the treatment of COPD which has also been shown to be effective over 24 hours after a single dose.(13)

About asthma and COPD

Asthma is a major, chronic airway disorder that is a serious public health problem in many

countries, and can have severe sometimes fatal consequences.(14) Asthma is one of the most common chronic diseases worldwide, affecting more than 300 million people(15) of whom an estimated 15 million suffer from severe disease.(16) Their health and quality of life are often severely affected, and more than 180,000 people worldwide are believed to die from asthma each year.(17)

COPD is also a major cause of chronic morbidity and mortality throughout the world, with many people dying prematurely from the disease or its complications.(18) COPD is currently the fourth leading cause of death in the world, and further increases in its prevalence can be predicted in the coming decades.(18) It is estimated that COPD is prevalent in 4% of the population in the USA, Europe and Japan,(19),(20),(21) and at least 15% of smokers will go on to develop the disease.(18) COPD, which encompasses chronic bronchitis and emphysema or both conditions, progresses slowly and eventually leads to a largely irreversible loss of lung function.

The foregoing press release contains forward-looking statements that can be identified by the use of forward-looking terminology such as could become, future therapeutic potential, should not, or similar expressions, or by express or implied discussions regarding the potential development and commercialization of indacaterol and NVA237. Such forward-looking statements involve known and unknown risks, uncertainties and other factors that may cause actual results to be materially different from any future results, performance or achievements expressed or implied by such statements. There can be no guarantee that the agreement that is the subject of this release will lead to commercialization of indacaterol or NVA237 in any market. Any such commercialization can be affected by, among other things, uncertainties relating to product development and clinical trials; regulatory actions or delays or government regulation generally; the ability to obtain or maintain patent or other proprietary intellectual property protection and competition in general; government, industry, and general public pricing pressures; as well as factors discussed in the Company's Form 20-F filed with the Securities and Exchange Commission. Should one or more of these risks or uncertainties materialize, or should underlying assumptions prove incorrect, actual results may vary materially from those described herein as anticipated, believed, estimated or expected. Novartis is providing this information as of this date and does not undertake any obligation to update any forward-looking statements contained in this document as a result of new information, future events or otherwise.

About Novartis

Novartis AG (NYSE: NVS) is a world leader in pharmaceuticals and consumer health. In 2004, the Group's businesses achieved net sales of USD 28.2 billion and pro forma net income of USD 5.6 billion. The Group invested approximately USD 4.2 billion in R&D. Headquartered in Basel, Switzerland, Novartis Group companies employ about 83,700 people and operate in over 140 countries around the world. For further information please consult <http://www.novartis.com>.

References

(1). Duvauchelle T, et al. Single-dose indacaterol, a novel once-daily 2-agonist, is well tolerated in patients with mild asthma. Poster (P1727) presented at 15th European Respiratory Society Congress, Copenhagen, Denmark, 17-21 September 2005.

(2). Beeh K-M, et al. Indacaterol: the first once-daily 2-agonist with 24-hour bronchodilation. Poster (P1725) presented at 15th European Respiratory Society Congress, Copenhagen, Denmark, 17-21 September 2005.

Edgar Filing: NOVARTIS AG - Form 6-K

(3). Tarral A, et al. Safety and tolerability of multiple-dose indacaterol, a novel once-daily 2-agonist, in patients with mild asthma. Poster (P1726) presented at 15th European Respiratory Society Congress, Copenhagen, Denmark, 17-21 September 2005.

(4). Chuchalin AG, et al. Cardiovascular safety of indacaterol, a novel once-daily 2-agonist, in patients with asthma. Poster (P1728) presented at 15th European Respiratory Society Congress, Copenhagen, Denmark, 17-21 September 2005.

(5). Kanniess F, et al. Indacaterol, a novel once-daily 2-agonist, demonstrates 24-hour efficacy and is well tolerated in patients with persistent asthma: a multiple-dose, dose-ranging study. Poster (P1729) presented at 15th European Respiratory Society Congress, Copenhagen, Denmark, 17-21 September 2005.

(6). Beier J, et al. Safety of multiple-dose indacaterol, a novel once-daily 2-agonist, in moderate-to-severe COPD. Poster (P1965) presented at 15th European Respiratory Society Congress, Copenhagen, Denmark, 17-21 September 2005.

Edgar Filing: NOVARTIS AG - Form 6-K

- (7). Aubier M, et al. Indacaterol, a novel once-daily 2-agonist, is effective and well tolerated on multiple dosing in patients with mild-to-moderate COPD. Poster (P1920) presented at 15th European Respiratory Society Congress, Copenhagen, Denmark, 17-21 September 2005.
- (8). Naline E, et al. Pharmacological characterization of indacaterol, a novel 2-agonist, on the isolated human bronchus. Poster (P1398) presented at 15th European Respiratory Society Congress, Copenhagen, Denmark, 17-21 September 2005.
- (9). Naline E, et al. Duration and onset of action of indacaterol, a novel once-daily 2-agonist, on the isolated human bronchus. Poster (P1399) presented at 15th European Respiratory Society Congress, Copenhagen, Denmark, 17-21 September 2005.
- (10). Lewis CA, et al. Indacaterol, a novel once-daily 2-agonist, demonstrates a long duration of action and fast onset *in vitro* and *in vivo* in the guinea pig. Poster (P835) presented at 15th European Respiratory Society Congress, Copenhagen, Denmark, 17-21 September 2005.
- (11). Battram CH, et al. Once-daily administration of indacaterol does not induce tachyphylaxis *in vivo*. Poster (P1400) presented at 15th European Respiratory Society Congress, Copenhagen, Denmark, 17-21 September 2005.
- (12). Lewis CA, et al. Indacaterol: preclinical evidence for an improved safety profile versus LABAs. Poster (P1401) presented at 15th European Respiratory Society Congress, Copenhagen, Denmark, 17-21 September 2005.
- (13). Langley SJ, et al. A randomized, double-blind, dose-ranging study to assess the effect of single doses of the inhaled anti-muscarinic AD 237 on FEV1 in patients with COPD. Proceedings of the American Thoracic Society 2005;2:A542.
- (14). Global Initiative for Asthma (GINA). NIH Publication 02-3659 issued January 1995 (updated 2004; accessed June 16, 2005). At: <http://www.ginasthma.com>
- (15). GINA. The Global Burden of Asthma Report 2004. At: <http://www.ginasthma.com>
- (16). American Thoracic Society. Proceedings of the ATS workshop on refractory asthma. Current understanding, recommendations, and unanswered questions. *Am J Respir Crit Care Med* 2000; 162:2341-2351.
- (17). World Health Organization. At: <http://www.who.int/mediacentre/factsheets/fs206/en/>

Edgar Filing: NOVARTIS AG - Form 6-K

(18). Global Initiative for Chronic Obstructive Lung Disease. Global strategy for the diagnosis, management and prevention of chronic obstructive pulmonary disease. Updated 2003; accessed June 16, 2005. At <http://www.goldcopd.com>

(19). National Institutes of Health National Health, Lung, and Blood Institute. Data Fact Sheet: Chronic Obstructive Pulmonary Disease. At: http://www.nhlbi.nih.gov/health/public/lung/other/copd_fact.pdf

(20). Gulsvik A. Mortality in and prevalence of chronic obstructive pulmonary disease in different parts of Europe. *Monaldi Arch Chest Dis*. 1999 Apr;54(2):160-2.

(21). Takemura H, Hida W, Sasaki T, Sugawara T, Sen T. Prevalence of Chronic Obstructive Pulmonary Disease in Japanese People on Medical Check-Up. *Tohoku J Exp Med* 2005;207(1):41-50.

###

© 2001 Novartis AG, Legal Disclaimer

Novartis International AG
Novartis Communications

CH-4002 Basel
Switzerland

Tel +41 61 324 2200
Fax + 41 61 324 3300
Internet Address:
<http://www.novartis.com>

MEDIA RELEASE COMMUNIQUE AUX MEDIAS MEDIENMITTEILUNG

Three-year study confirms long-term benefits of Xolair® for patients with severe allergic asthma

Basel, September 19, 2005 New long-term data show that Xolair® (omalizumab), a first-in-class monoclonal antibody for treating severe allergic asthma, helped patients to maintain control of their disease and was safe and well-tolerated in studies lasting more than three years. The data were presented today at the European Respiratory Society (ERS) congress in Denmark, along with results from seven clinical studies showing that Xolair significantly improved the lung function of patients.

Long-term control of severe allergic asthma was demonstrated in a study spanning 180 weeks of observation. The design included a 32-week randomised, double-blind, parallel-group, placebo-controlled study, followed by a 96-week open-label extension and a further 52-week extension.

As a result, patients completing the study and both extensions had been treated for more than three years. Disease control was maintained throughout the follow-up period in Xolair patients, who exhibited lower than expected changes in lung function and reduced use of inhaled corticosteroids (ICS).(1)

These results confirm that omalizumab potentially provides an important breakthrough in the fight against allergic asthma, said Prof. Marc Humbert of the Service de Pneumologie et Réanimation Respiratoire, Hôpital Antoine Bécclère, Clamart, France. Many people suffering from severe asthma have a substantially impaired quality of life and endure the constant fear that their next attack may prove fatal. By reducing exacerbations and the burden of disease, anti-IgE therapy offers a new approach to the treatment of this intractable condition.

Xolair manages asthma by targeting an underlying cause of allergic disease (up to 90% of asthma being allergic in origin).(2) It is designed to block the action of IgE, which is responsible for initiating the cascade of inflammatory symptoms such as airway constriction, mucus production, wheezing and shortness of breath. Xolair has been shown to decrease asthma exacerbations in some of the most difficult-to-treat patients whose condition remains inadequately-controlled despite the best conventional therapy, including inhaled corticosteroids, long-acting beta2-agonists and other controller medications.

Edgar Filing: NOVARTIS AG - Form 6-K

The Committee for Medicinal Products for Human Use (CHMP) recently gave a positive opinion on initial marketing authorisation for Xolair, opening the way for approval by the European Commission and for the therapy to become available in EU countries.

More about the three-year study demonstrating efficacy and safety

In the 32-week placebo controlled study, 341 patients received Xolair (at least 0.016 mg/kg/IgE [IU/mL] every four weeks) or placebo. A total of 222 patients entered Extension 1 (96 weeks), 178 of whom continued into Extension 2 (52 weeks). Patients in the extensions maintained stable lung function (measured by forced expiratory volume in one second, or FEV1). Mean FEV1 was 2.24 litres at the start of Extension 1 and 2.26 litres at the end of Extension 2, indicating persistency of therapeutic benefit from Xolair. In addition, more than 80% of patients had good/excellent asthma control during the extensions. Patients receiving inhaled corticosteroids at the start of Extension 1 and remaining on the same steroid reduced their ICS use by an average of 11% between the start of Extension 1 and the end of Extension 2.

In this long-term study, the overall incidence of adverse events was similar in the Xolair and placebo groups during the 32-week placebo-controlled study and Extension 1 (i.e. approximately 80% in both groups). During Extension 2, 134 patients (75%) had at least one adverse event, most of which were mild to moderate in severity. Serious adverse events remained infrequent during the second extension (4.5%) and were deemed by the treating physician to be unrelated to Xolair.(3)

Improvements in lung function and quality of life

Pooled data were also presented from seven trials in which allergic asthma patients (93% of whom had severe persistent disease) received Xolair as an add-on to inhaled corticosteroids, with or without oral corticosteroids and long-acting beta2-agonists. Out of 4,308 patients in the studies, FEV1 data were available for 3,537 patients (Xolair 2,443; control 1,094). Among Xolair recipients, 29.1% increased their FEV1 by ≥ 200 mL while 17.5% showed a ≥ 200 mL decrease. In the control group, 26.4% had a ≥ 200 mL increase in FEV1 and 26.1% had a ≥ 200 mL decrease. This equates to a net benefit of 11.6% for Xolair patients versus 0.3% for control (P<0.0001).(4)

Results from the same seven studies showed that add-on Xolair significantly improved patients' asthma-related quality of life, measured using the Juniper Adult Asthma Quality of Life Questionnaire (AQLQ). Quality of life data were available for 2,288 patients (Xolair 1,258; control 1,030). Those receiving Xolair had a mean increase of 1.01 points in their overall AQLQ score, compared to 0.67 in the control group (P<0.0001). Significantly more patients achieved a clinically meaningful improvement (i.e. an improved score of at least 0.5) in their AQLQ score from baseline in the Xolair group (66.3%) than the control group (53.2%, p<0.0001).(5)

Anti-IgE and severe asthma

Xolair has been available in the US for treating moderate to severe allergic asthma since July 2003, and as of 30 June 2005 it had been prescribed to more than 45,000 patients. In July 2005 the CHMP gave a positive opinion on Xolair in the EU, and approval is expected later this year. If approved by the European Commission, Xolair will be indicated as add-on therapy to improve asthma control in adult and adolescent patients (12 years of age and above) with severe persistent allergic asthma, who had the following, despite daily high-dose inhaled corticosteroids plus a long-acting inhaled beta2-agonist:

a positive skin test or *in vitro* reactivity to a perennial aeroallergen

reduced lung function (FEV1 <80%)

frequent daytime symptoms or night-time awakenings

multiple documented severe asthma exacerbations.

Xolair is also approved in Australia, Brazil, Canada, Dominican Republic, Guatemala, Israel, New Zealand and Venezuela. It was developed under an agreement between Novartis Pharma AG, Genentech, Inc., and Tanox, Inc.

Around 300 million people worldwide have asthma,(6) and an estimated 15 million of them suffer

from a severe form of the disease.(7) Their health and quality of daily life are often severely affected, and more than 180,000 people are believed to die from asthma each year throughout the world.(8)

The potential benefits of anti-IgE therapy are already recognised in treatment guidelines such as those developed by the Global Initiative for Asthma (GINA). These recommend anti-IgE therapy as add-on treatment for patients with severe allergic asthma that is inadequately controlled by standard clinical options.(9)

The foregoing release contains certain forward-looking statements that can be identified by terminology such as "potentially", "opening the way", "is expected", "will be", or similar expressions, or by express or implied discussions regarding the potential that Xolair will be approved for sale in any additional markets, or regarding any potential future revenues from Xolair. Such forward-looking statements involve known and unknown risks, uncertainties and other factors that may cause actual results with Xolair to be materially different from any future results, performance or achievements expressed or implied by such statements. There can be no guarantee that Xolair will be approved for sale in any additional market, or that it will achieve any particular sales level. In particular, management's expectations regarding commercialization of Xolair could be affected by, among other things, uncertainties relating to clinical trials; new clinical data, or additional analysis of existing clinical data; unexpected regulatory actions or delays or government regulation generally; the company's ability to obtain or maintain patent or other proprietary intellectual property protection; competition in general; government, industry and general public pricing pressures; as well as other risks and factors referred to in the Company's current Form 20-F on file with the US Securities and Exchange Commission. Should one or more of these risks or uncertainties materialize, or should underlying assumptions prove incorrect, actual results may vary materially from those anticipated, believed, estimated or expected. Novartis is providing the information in this press release as of this date and does not undertake any obligation to update any forward-looking statements contained in this press release as a result of new information, future events or otherwise.

About Novartis

Novartis AG (NYSE: NVS) is a world leader in pharmaceuticals and consumer health. In 2004, the Novartis Group's businesses achieved sales of USD 28.2 billion and pro forma net income of USD 5.6 billion. The Group invested approximately USD 4.2 billion in R&D. Headquartered in Basel, Switzerland, Novartis Group companies employ about 83,700 people and operate in over 140 countries around the world. For further information please consult <http://www.novartis.com>.

References

(1) Chung KF *et al.* Long-term asthma control with omalizumab, an anti-IgE monoclonal antibody in patients with severe allergic asthma. Poster (P417) presented at 15th European Respiratory Society Congress, Copenhagen, Denmark, 17-21 September 2005.

(2) Holt PG, Macaubas C, Stumbles PA, Sly PD. The role of allergy in the development of asthma. *Nature* 1999;402(Suppl.): B12-17.

(3) Chuchalin AG *et al.* Long-term safety and tolerability of omalizumab, an anti-IgE monoclonal antibody, in patients with severe allergic asthma. Poster (P421) presented at 15th European Respiratory Society Congress, Copenhagen, Denmark, 17-21 September 2005.

(4) Holgate S *et al.* Omalizumab significantly improves FEV1 in patients with severe persistent allergic asthma: a pooled analysis. Poster (P419) presented at 15th European Respiratory Society Congress, Copenhagen, Denmark, 17-21 September 2005.

(5) Beeh K-M *et al.* Omalizumab significantly improves quality of life in patients with severe persistent allergic asthma: a pooled analysis. Poster (P424) presented at 15th European Respiratory Society Congress, Copenhagen, Denmark, 17-21 September 2005.

(6) GINA. The Global Burden of Asthma Report 2004. <http://www.ginasthma.com>

(7) American Thoracic Society. Proceedings of the ATS workshop on refractory asthma. Current understanding, recommendations, and unanswered questions. *Am J Respir Crit Care Med* 2000; 162:2341-2351.

(8) World Health Organization. <http://www.who.int/mediacentre/factsheets/fs206/en/>

(9) Global Strategy for Asthma Management and Prevention 2004; 126-132. <http://www.ginasthma.com>

###

Novartis International AG
Novartis Communications

CH-4002 Basel
Switzerland

Tel +41 61 324 2200
Fax + 41 61 324 3300
Internet Address:
<http://www.novartis.com>

MEDIA RELEASE COMMUNIQUE AUX MEDIAS MEDIENMITTEILUNG

Novartis signs global licensing agreement to develop new topical applications of terbinafine the most prescribed oral treatment for fungal nail infections

Basel, September 15, 2005 Novartis announced today that it has signed a global licensing agreement with NexMed, Inc (NASDAQ: NEXM) to develop new topical forms of terbinafine, the active ingredient in its popular Lamisil® (terbinafine tablets) treatment for fungal nail infections.

Novartis has acquired the exclusive rights to further develop and commercialize NexMed's NM100060 nail lacquer treatment for onychomycosis, which is commonly known as nail fungus, and for any other indications.

Patients and doctors have said that they want a range of effective treatment options for onychomycosis. The currently available topical treatments for nail infections, while often preferred by patients, provide often poor efficacy, leading to patient frustration and poor compliance, said Joerg Reinhardt, Head of Global Development, Novartis Pharma AG. This agreement combines Lamisil's proven track record of effectively and safely treating people with onychomycosis with the potential of meeting the needs of patients who want an effective and simple-to-use topical option for the treatment of nail infections.

NM100060 is currently in Phase I development with the active ingredient terbinafine, which is also the active ingredient in Lamisil, the most frequently prescribed treatment for fungal nail infection worldwide. Under the terms of the agreement, Novartis has acquired the exclusive worldwide rights to develop and commercialize NM100060, which incorporates NexMed's NexACT® drug delivery technology to effectively enhance penetration of terbinafine in the infected nail areas.

More about Lamisil

Lamisil oral tablets are the No. 1 prescribed treatment based on sales for fungal nail infections with a proven efficacy and safety profile. More than 17 million people have treated their infections with Lamisil in the US since 1996.

Edgar Filing: NOVARTIS AG - Form 6-K

Lamisil contains the active ingredient terbinafine, which inhibits production of fungal cells by disrupting the fungal cell membranes. Terbinafine travels to the infected area under the toenail or fingernail since it is also attracted to keratin, which may also disrupt the infection. Because terbinafine accumulates and lingers in the area of the nail that is growing, it can continue to work for months even after the prescription is finished, allowing the nail to heal naturally over a period of time.

More about onychomycosis

Onychomycosis is a fungal infection affecting the toenails and/or fingernails. It is one of the most common dermatological diseases, affecting an estimated 30 million people in the US. Nail fungus can be painful, embarrassing, expensive and difficult to treat. A 1998 *American Journal of Dermatology* article estimated costs of monitoring and treatment of nail fungus ranged between USD 700 and USD 1,200 per patient annually.

The foregoing release contains forward-looking statements that can be identified by terminology such as "to develop and commercialize", "potentially meeting", or similar expressions, or by express or implied discussions regarding the potential development and commercialization of NM100060. Such forward-looking statements involve known and unknown risks, uncertainties and other factors that may cause actual results to be materially different from any future results, performance or achievements expressed or implied by such statements. There can be no guarantee that the agreement that is subject of this release will lead to commercialization of NM100060 in any market. In particular, management's expectations regarding commercialization of NM100060 could be affected by, among other things, uncertainties relating to product development and clinical trials; unexpected regulatory actions or delays or government regulation generally; the company's ability to obtain or maintain patent or other proprietary intellectual property protection; competition in general; industry, government, and general public pricing pressures; and other risks and factors referred to in Novartis AG's current Form 20-F on file with the US Securities and Exchange Commission. Should one or more of these risks or uncertainties materialize, or should underlying assumptions prove incorrect, actual results may vary materially from those anticipated, believed, estimated or expected. Novartis is providing the information in this press release as of this date and does not undertake any obligation to update any forward-looking statements contained in this press release as a result of new information, future events, or otherwise.

About Novartis

Novartis AG (NYSE: NVS) is a world leader in pharmaceuticals and consumer health. In 2004, the Novartis Group's businesses achieved sales of USD 28.2 billion and pro forma net income of USD 5.6 billion. The Group invested approximately USD 4.2 billion in R&D. Headquartered in Basel, Switzerland, Novartis Group companies employ about 83,700 people and operate in over 140 countries around the world. For further information please consult <http://www.novartis.com>.

###

© 2001 Novartis AG, Legal Disclaimer